



In re Appln. of KIMURA et al.  
Application No. 09/529,717



wherein  $R_3$  represents an alkyl group having 1 to 8 carbon atoms;  $n$  represents an integer of 1 to 4; and  $X$  represents an  $n$ -valent alcohol residue, having 1 to 18 carbon atoms, which optionally contains a hetero atom and/or a cyclic group,



wherein  $R_4$  represents an alkyl group having 1 to 8 carbon atoms;  $R_5$  and  $R_6$  independently represent a hydrogen atom or an alkyl group, having 1 to 18 carbon atoms, which optionally contains a hetero atom;  $m$  represents an integer of 1 to 3;  $Y$  represents an  $m$ -valent group, and when  $m$  is 1, it represents a hydrogen atom or an alkyl group, having 1 to 18 carbon atoms, which optionally contains a hetero atom, when  $m$  is 2, it represents a sulfur atom, an oxygen atom or an alkylidene group having 1 to 4 carbon atoms, and when  $m$  is 3, it

In re Appln. of KIMURA et al.  
Application No. 09/529,717

represents an isocyanuric acid- $N,N',N''$ -trimethylene group or a 1,3,5-trimethylbenzene-2,4,6-trimethylene group, and

(b) an amide represented by the following general formula (I):



wherein  $R_1$  represents an alkyl group having 12 to 21 carbon atoms, in a polyurethane.